

Molina Clinical Policy

Interspinous Decompression and Interlaminar Stabilization Devices for Spinal Stenosis

Policy No. 222



Last Approval: 04/08/2026
Next Review Due By: April 2027

DISCLAIMER

This Molina Clinical Policy (MCP) is intended to facilitate the Utilization Management process. Policies are not a supplementation or recommendation for treatment; Providers are solely responsible for the diagnosis, treatment, and clinical recommendations for the Member. It expresses Molina's determination as to whether certain services or supplies are medically necessary, experimental, investigational, or cosmetic for purposes of determining appropriateness of payment. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that this service or supply is covered (e.g., will be paid for by Molina) for a particular Member. The Member's benefit plan determines coverage – each benefit plan defines which services are covered, which are excluded, and which are subject to dollar caps or other limits. Members and their Providers will need to consult the Member's benefit plan to determine if there are any exclusion(s) or other benefit limitations applicable to this service or supply. If there is a discrepancy between this policy and a Member's plan of benefits, the benefits plan will govern. In addition, coverage may be mandated by applicable legal requirements of a State, the Federal government or CMS for Medicare and Medicaid Members. CMS's Coverage Database can be found on the CMS website. The coverage directive(s) and criteria from an existing National Coverage Determination (NCD) or Local Coverage Determination (LCD) will supersede the contents of this MCP and provide the directive for all Medicare members. References included were accurate at the time of policy approval and publication.

OVERVIEW

Spinal stenosis is a narrowing of the spinal canal that can compress the spinal cord and nerve roots, resulting in low back pain, neurogenic claudication, and functional limitation. The severity of symptoms may vary widely among individuals, ranging from mild activity-related discomfort to substantial impairment in walking tolerance and daily function. Symptoms typically worsen with lumbar extension and improve with flexion. Interspinous implants have been developed as minimally invasive options to offload neural structures by limiting extension and maintaining segmental space (Levin 2025). Interspinous implants fall into two distinct categories reflecting differences in regulatory indications, biomechanical function, and clinical application.

Interspinous decompression devices are minimally invasive implants designed to provide indirect decompression by limiting lumbar extension and increasing interspinous distance. These implants are inserted percutaneously between adjacent spinous processes to provide indirect decompression without direct laminectomy. By distracting the interspinous space and restricting painful lumbar extension, interspinous decompression implants enlarge canal and foraminal dimensions under load. Stand-alone devices serve as a primary intervention for moderate lumbar spinal stenosis in patients with flexion-relieved symptoms who have failed conservative therapy.

Interlaminar stabilization devices are used only as adjuncts to surgical decompression. These implants are placed midline between the laminae following laminectomy or laminotomy and are intended to provide dynamic stabilization while preserving some physiologic motion. FDA labeling specifies that interlaminar stabilization devices, such as Coflex, must be implanted after direct decompression at the treated level(s), differentiating them from stand-alone spacers in both mechanism and intended use.

Regulatory Information

The Coflex Interlaminar Stabilization device (Companion Spine) is a spinous process spacer/plate prosthesis and received premarket approval for treatment of 1- or 2-level lumbar spinal stenosis (LSS) from L1-L5, as an adjunct to surgical decompression, in skeletally mature patients with at least moderately impaired function, who experience flexion-relieve buttock, groin, leg pain after at least six months of non-operative treatment (FDA 2012).

The Superior Interspinous Spacer system (Boston Scientific) received FDA premarket approval in May 2015 for the stand-alone treatment of moderate LSS. The device is indicated for skeletally mature patients experiencing neurogenic intermittent claudication due to moderate degenerative stenosis, with or without Grade 1 spondylolisthesis, who have undergone at least six months of non-operative treatment (FDA 2015).

COVERAGE POLICY

Interspinous decompression and interlaminar stabilization devices (e.g., Coflex, Superior, and any other devices) are considered **experimental, investigational, and unproven** for any indication, due to insufficient clinical evidence of safety and efficacy in published peer-reviewed medical literature.

Molina Clinical Policy

Interspinous Decompression and Interlaminar Stabilization Devices for Spinal Stenosis

Policy No. 222



Last Approval: 04/08/2026
Next Review Due By: April 2027

DOCUMENTATION REQUIREMENTS: Molina Healthcare reserves the right to require that additional documentation be made available as part of its coverage determination; quality improvement; and fraud; waste and abuse prevention processes. Documentation required may include, but is not limited to, patient records, test results and credentials of the provider ordering or performing a drug or service. Molina Healthcare may deny reimbursement or take additional appropriate action if the documentation provided does not support the initial determination that the drugs or services were medically necessary, not investigational, or experimental, and otherwise within the scope of benefits afforded to the member, and/or the documentation demonstrates a pattern of billing or other practice that is inappropriate or excessive.

SUMMARY OF MEDICAL EVIDENCE

Systematic Reviews and Meta-Analyses

Han et al. (2024) conducted a systematic review and meta-analysis of five RCTs totaling 555 patients analyzing the safety and efficacy data of interspinous process devices in the treatment of lumbar spinal stenosis (LSS). The studies were analyzed to reveal no significant differences in Visual Analogue Scale (VAS) leg pain (SMD - 0.08, 95% CI - 0.32 to 0.15) and back pain (SMD 0.09, 95% CI - 0.27 to 0.45), Oswestry Disability Index (ODI) scores (MD 1.08, 95% CI - 11.23 to 13.39) and Zurich Claudication Questionnaire (ZCQ) physical function (MD - 0.09, 95% CI - 0.22 to 0.05) for interspinous process devices compared with decompression surgery. In terms of ZCQ symptom severity (MD - 0.22, 95% CI - 0.27 to - 0.16), decompression surgery showed superior to the interspinous process devices. As for complications (RR 1.08, 95% CI 0.36 to 3.27), the interspinous process devices had no advantages compared to decompression surgery and was inferior to it in reoperation rate (RR 2.58, 95% CI 1.67 to 3.96). The authors concluded there was no superiority in the clinical outcome for interspinous process devices compared with decompression surgery, and urged more clinical studies are warranted to determine the efficacy and safety of interspinous process devices.

Liang et al. (2022) published a systematic review and network meta-analysis that compared 20 RCTs that contained at least two of the following surgical procedures: (a) bilateral decompression via the unilateral approach, (b) decompression with conventional laminectomy, (c) decompression with fusion, (d) endoscopic decompression, (e) interspinous process devices only, (f) decompression with interlaminar stabilization, (g) decompression with lumbar spinal process-splitting laminectomy, and (h) minimally invasive tubular decompression. A total of 2201 patients were included in the meta-analysis. The primary outcomes of the meta-analysis were the ODI score, and secondary outcomes included VAS, SF-36, operation time, duration of hospital stay, reoperation, complications, and blood loss. Lower ODI scores were considered superior. The results of the meta-analysis found that decompression with interlaminar stabilization was significantly superior compared to bilateral decompression via the unilateral approach when comparing ODI scores. In terms of VAS scores, decompression with lumbar spinal process-splitting laminectomy was superior to decompression with conventional laminectomy, interspinous process devices, and unknown decompression (the procedure was uncertain or involved multiple options). The operation time of interspinous process devices was significantly shorter than all other options. However, interspinous process devices had a much higher rate of reoperation than other surgical methods of decompression. The most promising routine surgical option for most patients with LSS was the minimally invasive tubular decompression.

Tram et al. (2020) conducted a systematic review of decompression surgery versus interspinous devices for LSS. Twenty-five decompression-exclusive clinical trials totaling 3,386 patients and a mean age of 68.7 years, reported a 2.2% incidence rate of dural tears and a 2.6% incidence rate of postoperative infections. Eight interspinous devices exclusive clinical trials totaling 1,496 patients and a mean age of 65.1, reported a 5.3% incidence rate of postoperative leg pain and a 3.7% incidence rate of spinous process fractures. Seven studies that compared interspinous devices and decompression totaling 624 patients, found a reoperation rate of 8.3% in interspinous devices patients vs. 3.9% in decompression patients and dural tears in 0.32% of interspinous devices patients vs. 5.2% in decompression patients. Utilizing a random-effects model the difference between preoperative and the 1-2-year postoperative VAS scores between interspinous device surgery and lumbar decompression was analyzed to reveal no difference between the groups. The authors concluded that decompression and interspinous devices are unique surgical interventions with different therapeutic efficacies and complications. The collected studies do not consistently demonstrate superiority of either procedure over the other.

Molina Clinical Policy

Interspinous Decompression and Interlaminar Stabilization Devices for Spinal Stenosis

Policy No. 222



Last Approval: 04/08/2026
Next Review Due By: April 2027

Coflex

Systematic Reviews and Meta-Analyses

Fan and Zhu (2020) conducted a network meta-analysis of studies comparing decompression alone versus fusion and Coflex in the treatment of lumbar degenerative disease. A total of 10 randomized controlled trials were included totaling 946 patients. Compared with decompression alone group, there were no significant differences of ODI in Coflex and lumbar interbody fusion groups after surgery. Coflex and posterior lumbar interbody fusion were better in decreasing VAS score compared with decompression alone. In addition, Coflex have a less complication incidence rate. A total analysis of the data led the authors to conclude that while the effectiveness and safety of the fusion and Coflex techniques are still not clear, Coflex and lumbar interbody fusion had the similar effectiveness in improving lumbar function and quality of life, with the added benefit of increase in pain relief reporting and a lower complication incidence rate.

Non-Randomized Studies, Retrospective Reviews and Other Evidence

Hayes (2024) published an update to the Health Technology Assessment report on the use of the CoFlex Interlaminar Stabilization device for the treatment of lumbar spinal stenosis in adults. The available evidence, although of low quality, indicates that outcomes for the CoFlex device combined with decompression are comparable to decompression with fusion over an 8-year period, and to decompression alone over a 2-year period. Adverse event rates appeared to be similar across groups using the CoFlex device and its comparators. Additionally, the CoFlex device may offer advantages in terms of reduced operative time and shorter hospital stays. Hayes emphasized that the uncertainty surrounding the evidence is largely due to the fact that all studies reviewed were of low quality, providing limited insight into the long-term benefits of the CoFlex device compared to traditional surgical approaches. Additionally, the absence of clear criteria for patient selection further contributes to this uncertainty.

Davis et al. (2013a) compared the efficacy and safety of spinal decompression plus Coflex with decompression plus fusion in 322 patients with LSS, and in a subset of 150 patients with grade I spondylolisthesis. Both treatments led to significant improvement at 24 months in mean scores on the VAS for back pain and leg pain, ODI, SF-12 physical component, and ZCQ symptom severity and physical function, compared with baseline values. At 24 months, mean scores for the SF-12 physical component and ZCQ symptom severity, physical function, and patient satisfaction were significantly better for the Coflex than for fusion; however, mean VAS and ODI scores were similar for the 2 approaches in the entire cohort. In the entire cohort and the in the subset with spondylolisthesis, the mean SF-12 mental component score did not change appreciably and was similar between the Coflex and Fusion groups at all evaluation times. At 24 months, radiographic results revealed changes in ROM in patients who had fusion (rotation and translation decreased at the treated lumbar level(s) and increased at the level above and the level below the treated level(s)). In contrast, ROM was fairly well preserved (rotation and translation changed by $< 1.0^\circ$ or < 1.0 mm, respectively, at treated and adjacent levels) in the Coflex group.

Bae et al. (2016) performed a three-year follow-up analysis of the Davis (2013a) RCT. At 36 months, 91% (195/215) of the Coflex group and 88% (94/107) of the fusion group were included in the analysis. The initial efficacy endpoints (composite scores) were modified for use at 36 months. At 36 months, 62.2% of the individuals in the Coflex group compared to 48.9% of the individuals in the 94-group reported composite clinical success scores (difference = 13.3%, 95% confidence interval [CI]; 1.1%-25.5%, $p=0.03$). There are several limitations in this study including the limited follow-up period and the heterogeneous mix of individuals including those without spondylolisthesis for which fusion/stabilization is an unproven procedure

Davis et al. (2013b) reported on the outcomes of a subset of 150 patients with Meyerding grade I spondylolisthesis ($\leq 25\%$ sagittal plane translation on flexion-extension radiographs) who were included in the randomized FDA IDE trial in the Coflex group ($n=99$; mean age 63.1 years, range not reported; 41% men; 2-level procedures required in 64.2%; mean ODI 59.4; mean VAS for back pain 80.3; mean VAS for worse leg pain 77.9) or the Fusion group ($n=51$; mean age 65.0 years, range not reported; 19% men; 2-level procedures required in 63.6%; mean ODI 60.0; mean VAS for back pain 78.6; mean VAS for worse leg pain 79.1). Follow-up findings were reported only for the 24-month evaluation, at which time data were available in 94.9% of the Coflex group and 94.1% of the Fusion group. Both groups demonstrated significant improvement in mean scores for ODI (-38.3 and -37.1 points, respectively), VAS for back pain (-54.9 and -58.0 mm), VAS for worse leg pain (-58.9 and -56.2 points), SF-12 physical component (16.4 and 14.8 points), ZCQ symptom severity (-1.64 and -1.40 points), and ZCQ physical function (-1.24 and -1.10 points). The rate of composite clinical success was similar in the Coflex and Fusion groups (62.8% and 62.5%, respectively).

Molina Clinical Policy

Interspinous Decompression and Interlaminar Stabilization Devices for Spinal Stenosis

Policy No. 222



Last Approval: 04/08/2026
Next Review Due By: April 2027

Superion

Randomized Controlled Trials

Nunley et al. (2017a) reported five-year clinical outcomes of a randomized controlled U.S. FDA noninferiority trial in individuals with moderate lumbar spinal stenosis. While the original trial compared the Superion to the X STOP device, the analysis was restricted to the Superion trial arm. A total of 73% (88/121) of the living individuals who received the spacer device participated in the 5-year clinical outcomes assessment. Outcomes were assessed using the ZCQ, leg and back pain severity by VAS, and the ODI. The authors reported success rates in all areas of assessment, 84% reported clinical success in at least two of the three ZCQ domains, 80% leg pain VAS scores, 65% back pain VAS scores and 65% for ODI scores. There remains a lack of studies which compare interspinous spacers to standard treatments, such as decompression surgery.

Non-Randomized Studies, Retrospective Reviews and Other Evidence

Hayes (2024) published an update to the Evolving Evidence Review on the Superion interspinous spacer (ISS). The review evaluated clinical studies, systematic reviews (none of which met inclusion criteria), and guidelines for the treatment of lumbar spinal stenosis and neurogenic claudication using the Superion device. The assessed studies were determined to be of very poor or poor quality, and no comparative research was identified. Hayes concluded that there is no evidence to suggest that the Superion ISS offers advantages over other treatment options, such as fusion surgery or existing commercial alternatives. An analysis of guidelines and position statements revealed inconsistent support for the Superion ISS, specifically in its role in addressing lumbar spinal stenosis with neurogenic claudication. While some guidelines deemed the evidence sufficient to endorse its use, others found the evidence insufficient. Consequently, the long-term health outcomes associated with the Superion ISS remain uncertain, underscoring the need for further investigation.

Hagedorn et al. (2022) conducted a retrospective study to assess the rate of lumbar decompression surgery following minimally invasive procedures, specifically the MILD (Minimally Invasive Lumbar Decompression) procedure and the placement of the Superion Indirect Decompression device. The study included patients with lumbar spinal stenosis (LSS) who underwent MILD and/or Superion between January 2011 and July 2019. Eligible participants had a follow-up of at least two years, pre-procedural MRI findings, and surgical records. The final analysis comprised 199 patients: 28.6% underwent only MILD, 62.3% underwent only Superion, and 9.0% received MILD followed by Superion. Notably, two patients underwent the MILD procedure twice at the same spinal level. Over the two-year follow-up period, 2.0% of patients (four individuals) required subsequent lumbar surgery, with rates of 5.3% for MILD and 0.8% for Superion. The study acknowledged that some patients might not have been candidates for surgery, which could have influenced the results. The authors concluded that minimally invasive decompression treatments for LSS are associated with low rates of follow-up surgeries, potentially resulting in cost savings and a reduction in severe adverse events (AEs). The low surgical rates may be attributed to symptom improvement, patient preferences to avoid surgery, or the patients being deemed unsuitable for surgical intervention. It should be noted that the lead author is a consultant for Boston Scientific (device manufacturer) and disclosed this conflict of interest.

Patel et al. (2015b) published a report on 3-year durability of results of the pivotal trial. At 36 months, the overall treatment success (primary composite endpoint) remained stable in the Superion group (52.5% of 120 participants available for follow-up at 36 months versus 52.7% at 24 months). In the X-Stop group, the composite endpoint of overall treatment success was 38.0% of 129 participants available for follow-up at 36 months, reduced from 50.2% at 24 months. The difference between groups was statistically significant ($P=0.023$). A total of 26 (14%) participants in the Superion group required surgical decompression within 3 years. Most patients in the Superion group experienced significant improvements in individual outcome measures, including back pain as measured by a ≥ 20 mm decrease in VAS (76.8%), VAS leg pain (84.1%), ZCQ physical function (80.5%), ZCQ symptom severity (82.9%), ODI (≥ 15 -point decrease) (69.5%), and ZCQ patient satisfaction (91.5%) at 36 months. Between-group differences in most individual outcome measures were not statistically significant, except for VAS leg pain. A total of 69.7% of patients in the X-Stop group had durable improvement in leg pain at 36 months, compared with 84.1% of the Superion group ($P=0.037$).

National and Specialty Organizations

The **North American Spine Society** (2025) published coverage policy recommendations for lumbar interspinous devices with decompression. These recommendations pertain to all interspinous process devices intended for

Molina Clinical Policy
Interspinous Decompression and Interlaminar Stabilization
Devices for Spinal Stenosis
Policy No. 222



Last Approval: 04/08/2026
 Next Review Due By: April 2027

placement during the same surgery as direct decompression procedures. Interspinous distraction devices without fusion may be considered appropriate for degenerative lumbar stenosis, provided specific criteria are met. These criteria include the presence of neurogenic claudication that improves with lumbar flexion, patients being over 50 years old, a failure of nonoperative treatment, no more than 25° of degenerative scoliosis, and no greater than a Grade I degenerative spondylolisthesis.

The **International Society for the Advancement of Spinal Surgery** views interlaminar stabilization following direct decompression as a nonfusion alternative that could offer greater stability than decompression alone (Guyer et al., 2016). They regard the evidence from studies comparing decompression alone to decompression combined with interlaminar stabilization using the Coflex device as persuasive in supporting this treatment for carefully selected patients. Eligible patients for decompression with interlaminar stabilization include radiographically confirmed moderate lumbar spinal stenosis (LSS) at levels L1 to L5 requiring decompression, no instability at the index or adjacent spinal levels, pain relief with lumbar flexion (with or without associated back pain), and at least 12 weeks of attempted conservative management.

The **National Institute for Health and Clinical Excellence (NICE)** (2010) issued guidance on interspinous distraction procedures for lumbar spinal stenosis causing neurogenic claudication in 2010 which state that current evidence shows the procedures are, “efficacious for carefully selected patients in the short and medium term, although failure may occur, and further surgery may be needed. There are no major safety concerns; these procedures may be used provided that normal arrangements are in place for clinical governance, consent, and audit.”

CODING & BILLING INFORMATION

CPT (Current Procedural Terminology)

Code	Description
22867	Insertion of interlaminar/interspinous process stabilization/distraction device, without fusion, including image guidance when performed, with open decompression, lumbar; single level
22868	Insertion of interlaminar/interspinous process stabilization/distraction device, without fusion, including image guidance when performed, with open decompression, lumbar; second level (List separately in addition to code for primary procedure)
22869	Insertion of interlaminar/interspinous process stabilization/distraction device, without open decompression or fusion, including image guidance when performed, lumbar; single level
22870	Insertion of interlaminar/interspinous process stabilization/distraction device, without open decompression or fusion, including image guidance when performed, lumbar; second level (List separately in addition to code for primary procedure)

HCPCS (Healthcare Common Procedure Coding System)

Code	Description
C1821	Interspinous process distraction device (implantable)

CODING DISCLAIMER: Codes listed in this policy are for reference purposes only and may not be all-inclusive. Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement. Listing of a service or device code in this policy does not guarantee coverage. Coverage is determined by the benefit document. Molina adheres to Current Procedural Terminology (CPT®), a registered trademark of the American Medical Association (AMA). All CPT codes and descriptions are copyrighted by the AMA; this information is included for informational purposes only. Providers and facilities are expected to utilize industry standard coding practices for all submissions. When improper billing and coding is not followed, Molina has the right to reject/deny the claim and recover claim payment(s). Due to changing industry practices, Molina reserves the right to revise this policy as needed.

APPROVAL HISTORY

04/08/2026	Policy reviewed. No changes to coverage criteria. Updated Summary of Medical Evidence and References. Updated title to “Interspinous Decompression and Interlaminar Stabilization Devices for Spinal Stenosis”.
04/09/2025	Policy reviewed. No changes to coverage criteria. Updated Summary of Medical Evidence and References.
04/10/2024	Policy reviewed, no changes to coverage criteria. IRO Peer Review on February 27, 2024, by a practicing physician board certified in Orthopedic Surgery.

Molina Clinical Policy

Interspinous Decompression and Interlaminar Stabilization Devices for Spinal Stenosis

Policy No. 222



Last Approval: 04/08/2026
Next Review Due By: April 2027

04/13/2023	Policy reviewed, no changes to coverage statement. Updated references and Summary of Evidence. Added topping-off procedure to Overview.
04/13/2022	Policy reviewed, no changes to coverage statement. Updated references and Summary of Evidence.
04/05/2021	Policy reviewed, no changes. A review of clinical studies and guidelines suggests minimal support for using interspinous spacers for the treatment of lumbar spinal stenosis with neurogenic claudication.
04/23/2020	Policy reviewed, no changes.
06/19/2019	Policy reviewed, no changes, updated professional society guidelines and references.
07/10/2018	Policy reviewed, no changes to coverage. Added new device (Vertiflex's Superior® interspinous spacer system), updated Summary of Medical Evidence, references, and coding. IRO reviewed on March 27, 2018, by a practicing, board-certified physician in Orthopedic Surgery.
06/22/2017	Policy reviewed, no changes.
09/15/2016	Policy reviewed, no changes.
12/16/2015	Policy reviewed, no changes.
03/16/2015	New policy.

REFERENCES

1. Bae HW, Davis RJ, Laurysen C, Leary S, Maislin G, Musacchio M Jr. Three-year follow-up of the prospective, randomized, controlled trial of Coflex interlaminar stabilization vs instrumented fusion in patients with lumbar stenosis. *Neurosurgery*. 2016. doi: 10.1227/NEU.0000000000001237.
2. Chiou K, Chiu YC, Lee CY, Huang TJ, Lai YC, Yang CJ, Hsu JC, Wu MH. Comparison of long-term outcomes of spinal fusion surgeries supplemented with "topping-off" implants in lumbar degenerative diseases: A systematic review and network meta-analysis. *N Am Spine Soc J*. 2022 Oct 22; 12:100177. doi: 10.1016/j.xnsj.2022.100177. PMID: 36394053. PMCID: PMC9650073.
3. Chou PH, Lin HH, An HS, Liu KY, Su WR, Lin CL. Could the topping-off technique be the preventive strategy against adjacent segment disease after pedicle screw-based fusion in lumbar degenerative diseases? A systematic review. *Biomed Res Int*. 2017; 2017:4385620. doi: 10.1155/2017/4385620. Epub 2017 Feb 22. PMID: 28321409; PMCID: PMC5340959.
4. Davis RJ, Errico TJ, Bae H, Auerbach JD. Decompression and Coflex interlaminar stabilization compared with decompression and instrumented spinal fusion for spinal stenosis and low-grade degenerative spondylolisthesis. two-year results from the prospective, randomized, multicenter, Food and Drug Administration Investigational Device Exemption trial. *Spine (Phila Pa 1976)*. 2013a Aug 15;38(18):1529-39. doi: 10.1097/BRS.0b013e31829a6d0a.
5. Davis R, Auerbach JD, Bae H, Errico TJ. Can low-grade spondylolisthesis be effectively treated by either Coflex interlaminar stabilization or laminectomy and posterior spinal fusion? Two-year clinical and radiographic results from the randomized, prospective, multicenter US investigational device exemption trial: clinical article. *J Neurosurg Spine*. 2013b Aug;19(2):174-84. doi: 10.3171/2013.4.SPINE12636.
6. Fan Y, Zhu L. Decompression alone versus fusion and Coflex in the treatment of lumbar degenerative disease: A network meta-analysis. *Medicine (Baltimore)*. 2020 Mar;99(11): e19457. doi: 10.1097/MD.00000000000019457. PMID: 32176077; PMCID: PMC7220096.
7. Guyer R, Musacchio M, et al. ISASS recommendations/coverage criteria for decompression with interlaminar stabilization-coverage indications, limitations, and/or medical necessity. *Int J Spine Surg*. 2016 Dec 5; 10:41. doi: 10.14444/3041.
8. Han B, Chen Y, Liang W, Yang Y, Ding Z, Yin P, Hai Y. Is the interspinous process device safe and effective in elderly patients with lumbar degeneration? A systematic review and meta-analysis of randomized controlled trials. *Eur Spine J*. 2024 Feb 12. doi: 10.1007/s00586-023-08119-z. PMID: 38342843.
9. Hayes. Coflex interlaminar stabilization device (Surgalign Spine Technologies Inc.) for treatment of lumbar spinal stenosis. Health Technology Assessment. Published October 21, 2021. Updated October 30, 2024. Accessed February 12, 2026. <https://evidence.hayesinc.com/>.
10. Hayes. Superior Interspinous Spacer System (Vertiflex) for treatment of neurogenic claudication caused by spinal stenosis. Evolving Evidence Review. Published September 17, 2021. Reviewed October 17, 2024. Accessed February 12, 2026. <https://evidence.hayesinc.com/>.
11. Levin, K Lumbar spinal stenosis: Treatment and prognosis. Updated February 11, 2026. Accessed February 12, 2026. <http://www.uptodate.com>
12. Liang Z, Xu X, Chen X, Zhuang Y, Wang R, Chen C. Clinical evaluation of surgery for single-segment lumbar spinal stenosis: A systematic review and bayesian network meta-analysis. *Orthop Surg*. 2022 Jul;14(7):1281-1293. doi: 10.1111/os.13269. PMID: 35582931. PMCID: PMC9251271.
13. National Institute for Health and Care Excellence. Interspinous distraction procedures for lumbar spinal stenosis causing neurogenic claudication [IPG365]. Published: 24 November 2010. Accessed February 12, 2026. [NICE.org.uk](https://www.nice.org.uk).
14. North American Spine Society (NASS). Coverage recommendations: Interspinous device with decompression. Revised March 2025. Accessed February 12, 2026.
15. Nunley PD, Patel VV, Orndorff DG, et al. Five-year durability of stand-alone interspinous process decompression for lumbar spinal stenosis. *Clin Interv Aging*. 2017a Sep 6; 12:1409-1417. doi: 10.2147/CIA.S143503.
16. Patel VV, Nunley PD, Whang PG et al. Superior (®) InterSpinous Spacer for treatment of moderate degenerative lumbar spinal stenosis: Durable three-year results of a randomized controlled trial. *J Pain Res*. 2015b Oct 3; 8:657-62. doi: 10.2147/JPR.S92633.
17. Tram J, Srinivas S, Wali AR, Lewis CS, Pham MH. Decompression Surgery versus Interspinous Devices for Lumbar Spinal Stenosis: A Systematic Review of the Literature. *Asian Spine J*. 2020 Aug;14(4):526-542. doi: 10.31616/asj.2019.0105. PMID: 31906617; PMCID: PMC7435320.
18. United States Food and Drug Administration (FDA). Pre-market approval (PMA): X-Stop LTOS (multiple approvals). Product code NQO. PMA: P040001. [FDA.gov](https://www.fda.gov). Accessed February 12, 2026.
19. United States Food and Drug Administration (FDA). Pre-market approval (PMA): coflex Interlaminar Stabilization Device (multiple approvals). Product code NQO. PMA: P110008. [FDA.gov](https://www.fda.gov). Accessed February 12, 2026.
20. United States Food and Drug Administration (FDA). Pre-market approval (PMA): Superior Indirect Decompression System (IDS) (multiple approvals). Product code NQO. PMA: P140004. [FDA.gov](https://www.fda.gov). Accessed February 12, 2026.