

## POLICY SECTIONS

[DISCLAIMER](#) | [POLICY DESCRIPTION](#) | [RELATED POLICIES](#) | [INDICATIONS AND/OR LIMITATIONS OF COVERAGE](#) | [EXCLUSION CRITERIA](#) | [MEDICATION MANAGEMENT](#) | [ATTACHMENTS](#) | [APPLICABLE CPT / HCPCS PROCEDURE CODES](#) | [APPROVAL HISTORY](#) | [REFERENCES](#) | [APPENDIX](#)

## DISCLAIMER

This Molina Clinical Policy (MCP) is intended to facilitate the Utilization Management process. Policies are not a supplementation or recommendation for treatment; Providers are solely responsible for the diagnosis, treatment, and clinical recommendations for the Member. It expresses Molina's determination as to whether certain services or supplies are medically necessary, experimental, investigational, or cosmetic for purposes of determining appropriateness of payment. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that this service or supply is covered (e.g., will be paid for by Molina) for a particular Member. The Member's benefit plan determines coverage – each benefit plan defines which services are covered, which are excluded, and which are subject to dollar caps or other limits. Members and their Providers will need to consult the Member's benefit plan to determine if there are any exclusion(s) or other benefit limitations applicable to this service or supply. If there is a discrepancy between this policy and a member's plan of benefits, the benefits plan will govern. In addition, coverage may be mandated by applicable legal requirements of a State, the Federal government or CMS for Medicare and Medicaid Members. CMS's Coverage Database can be found on the CMS website. The coverage directive(s) and criteria from an existing National Coverage Determination (NCD) or Local Coverage Determination (LCD) will supersede the contents of this MCP and provide the directive for all Medicare members. References included were accurate at the time of policy approval and publication.

## POLICY DESCRIPTION

To define and describe the accepted indications for Kymriah (tisagenlecleucel) usage in the treatment of cancer, including FDA approved indications, and off-label indications.

The use of this drug must be supported by one of the following: FDA approved product labeling, CMS-approved compendia, National Comprehensive Cancer Network (NCCN), American Society of Clinical Oncology (ASCO) clinical guidelines, or peer-reviewed literature that meets the requirements of the CMS Medicare Benefit Policy Manual Chapter 15.

## INDICATIONS and/or LIMITATIONS OF COVERAGE

### **A. Continuation requests for a not – approvable medication shall be exempt from this policy provided:**

1. The requested medication was used within the last year; **AND**
2. The member has not experienced disease progression and/or no intolerance to the requested medication; **AND**
3. Additional medication(s) are not being added to the continuation request.

### **B. Acute Lymphoblastic Leukemia (ALL)**

1. Kymriah (tisagenlecleucel) is being used when the following criteria are met:
  - a. Member is 25 years old or younger, and has ALL with confirmed documentation of CD19 tumor expression (demonstrated in bone marrow or peripheral blood by flow cytometry); **AND**
  - b. Member has B- Cell ALL that is refractory or in second or later relapse

### **C. B-Cell Lymphomas**

1. Kymriah (tisagenlecleucel) may be used for members who are 18 years of age or older with relapsed or refractory large B-cell lymphoma after two or more lines of systemic therapy, including diffuse large B-cell lymphoma (DLBCL) not otherwise specified, high grade B-cell lymphoma, and DLBCL arising from follicular lymphoma

## EXCLUSION CRITERIA

- A. Kymriah (tisagenlecleucel) is being used after disease progression on or after CAR-T cell therapy directed towards CD19 antigen [Kymriah ((tisagenlecleucel), Breyanzi (lisocabtagene maraleucel), or Yescarta (axicabtagene ciloleucel)].
- B. CD-19 positivity not confirmed and documented.
- C. Active CNS involvement with lymphoma.
- D. Dosing exceeds single dose limit of Kymriah (tisagenlecleucel)  $6.0 \times 10^8$  CAR-positive viable T cells (for B-Cell Lymphomas);  $2.5 \times 10^8$  CAR-positive viable T cells (for ALL).
- E. Exceeds duration limit as one time administration.
- F. Investigational use of Kymriah (tisagenlecleucel) with an off-label indication that is not sufficient in evidence or is not generally accepted by the medical community. Sufficient evidence that is not supported by CMS recognized compendia or acceptable peer reviewed literature is defined as any of the following:
  1. Whether the clinical characteristics of the patient and the cancer are adequately represented in the published evidence.
  2. Whether the administered chemotherapy/biologic therapy/immune therapy/targeted therapy/other oncologic therapy regimen is adequately represented in the published evidence.
  3. Whether the reported study outcomes represent clinically meaningful outcomes experienced by patients. Generally, the definition of Clinically Meaningful outcomes are those recommended by ASCO, e.g., Hazard Ratio of  $< 0.80$  and the recommended survival benefit for OS and PFS should be at least 3 months.
  4. Whether the experimental design, in light of the drugs and conditions under investigation, is appropriate to address the investigative question. (For example, in some clinical studies, it may be unnecessary or not feasible to use randomization, double blind trials, placebos, or crossover).
  5. That non-randomized clinical trials with a significant number of subjects may be a basis for supportive clinical evidence for determining accepted uses of drugs.
  6. That case reports are generally considered uncontrolled and anecdotal information and do not provide adequate supportive clinical evidence for determining accepted uses of drugs.
  7. That abstracts (including meeting abstracts) without the full article from the approved peer- reviewed journals lack supporting clinical evidence for determining accepted uses of drugs.

## MEDICATION MANAGEMENT

Please refer to the FDA label/package insert for details regarding these topics.

## APPLICABLE CPT / HCPCS PROCEDURE CODES

### CPT (Current Procedural Terminology) Codes

Code	Description
<b>0537T</b>	Chimeric antigen receptor T-cell (CAR-T) therapy; harvesting of blood-derived T lymphocytes for development of genetically modified autologous CAR-T cells, per day
<b>0538T</b>	Chimeric antigen receptor T-cell (CAR-T) therapy; preparation of blood-derived T lymphocytes for transportation (e.g., cryopreservation, storage)
<b>0539T</b>	Chimeric antigen receptor T-cell (CAR-T) therapy; receipt and preparation of CAR-T cells for administration
<b>0540T</b>	Chimeric antigen receptor T-cell (CAR-T) therapy; CAR-T cell administration, autologous

### HCPCS (Healthcare Common Procedure Coding System) Code

Code	Description
Q2042	Tisagenlecleucel, up to 600 million CAR-positive viable T cells, including leukapheresis and dose preparation procedures, per therapeutic dose

**AVAILABLE DOSAGE FORMS:** Single-dose unit infusion bag: frozen suspension of genetically modified autologous T cells labeled for the specific recipient

**CODING DISCLAIMER.** Codes listed in this policy are for reference purposes only and may not be all-inclusive. Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement. Listing of a service or device code in this policy does not guarantee coverage. Coverage is determined by the benefit document. Molina adheres to Current Procedural Terminology (CPT®), a registered trademark of the American Medical Association (AMA). All CPT codes and descriptions are copyrighted by the AMA; this information is included for informational purposes only. Providers and facilities are expected to utilize industry standard coding practices for all submissions. When improper billing and coding is not followed, Molina has the right to reject/deny the claim and recover claim payment(s). Due to changing industry practices, Molina reserves the right to revise this policy as needed.

### APPROVAL HISTORY

04/10/2024	Policy reviewed. Changes to coverage criteria include: removal of relapse after allo HSCT under ALL indication, and specific stipulations for relapse under B-cell lymphoma indication. Removed bone marrow reserve, adequate renal/hepatic/cardiac/pulm function, hx of CNS disorder, hx of autoimmune disease, active infection, and previous allo HSCT from exclusion criteria.
06/14/2023	Policy revision to include members with relapsed/refractory ALL progressed after 2 lines of standard chemotherapy and added exclusion criteria-must have documented CD19. Removed codes C9399, J3490, J3590, and J9999.
08/10/2022	Adopted NCH policy and retired MCP.

### REFERENCES

1. Ellis LM, Bernstein DS, Voest EE, Berlin JD, Sargent D, Cortazar P, Garrett-Mayer E, Herbst RS, Lilenbaum RC, Sima C, Venook AP, Gonen M, Schilsky RL, Meropol NJ, Schnipper LE. American Society of Clinical Oncology perspective: Raising the bar for clinical trials by defining clinically meaningful outcomes. *J Clin Oncol*. 2014 Apr 20;32(12):1277-80. doi: 10.1200/JCO.2013.53.8009. Epub 2014 Mar 17. PMID: 24638016.
2. Fowler NH, Dickinson M, Dreyling M, Martinez-Lopez J, Kolstad A, Butler J, Ghosh M, Popplewell L, Chavez JC, Bachy E, Kato K, Harigae H, Kersten MJ, Andreadis C, Riedell PA, Ho PJ, Pérez-Simón JA, Chen AI, Nastoupil LJ, von Tresckow B, Ferreri AJM, Teshima T, Patten PEM, McGuirk JP, Petzer AL, Offner F, Viardot A, Zinzani PL, Malladi R, Zia A, Awasthi R, Masood A, Anak O, Schuster SJ, Thieblemont C. Tisagenlecleucel in adult relapsed or refractory follicular lymphoma: the phase 2 ELARA trial. *Nat Med*. 2022 Feb;28(2):325-332. doi: 10.1038/s41591-021-01622-0. Epub 2021 Dec 17. PMID: 34921238.
3. Kymriah prescribing information. Novartis Pharmaceuticals Corporation. East Hanover, NJ 2022.
4. Maude SL, Frey N, Shaw PA, Aplenc R, Barrett DM, Bunin NJ, Chew A, Gonzalez VE, Zheng Z, Lacey SF, Mahnke YD, Melenhorst JJ, Rheingold SR, Shen A, Teachey DT, Levine BL, June CH, Porter DL, Grupp SA. Chimeric antigen receptor T cells for sustained remissions in leukemia. *N Engl J Med*. 2014 Oct 16;371(16):1507-17. doi: 10.1056/NEJMoa1407222. Erratum in: *N Engl J Med*. 2016 Mar 10;374(10):998. PMID: 25317870; PMCID: PMC4267531.
5. Medicare Benefit Policy Manual Chapter 15 Covered Medical and Other Health Services. Updated March 16, 2023. Accessed June 7, 2023. <https://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/Downloads/bp102c15.pdf>.
6. Schuster SJ, Bishop MR, Tam CS, Waller EK, Borchmann P, McGuirk JP, Jäger U, Jaglowski S, Andreadis C, Westin JR, Fleury I, Bachanova V, Foley SR, Ho PJ, Mielke S, Magenau JM, Holte H, Pantano S, Pacaud LB, Awasthi R, Chu J, Anak Ö, Salles G, Maziarz RT; JULIET Investigators. Tisagenlecleucel in Adult Relapsed or Refractory Diffuse Large B-Cell Lymphoma. *N Engl J Med*. 2019 Jan 3;380(1):45-56. doi: 10.1056/NEJMoa1804980. Epub 2018 Dec 1. PMID: 30501490.