

## DISCLAIMER

This Molina Clinical Policy (MCP) is intended to facilitate the Utilization Management process. Policies are not a supplementation or recommendation for treatment; Providers are solely responsible for the diagnosis, treatment, and clinical recommendations for the Member. It expresses Molina's determination as to whether certain services or supplies are medically necessary, experimental, investigational, or cosmetic for purposes of determining appropriateness of payment. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that this service or supply is covered (e.g., will be paid for by Molina) for a particular Member. The Member's benefit plan determines coverage – each benefit plan defines which services are covered, which are excluded, and which are subject to dollar caps or other limits. Members and their Providers will need to consult the Member's benefit plan to determine if there are any exclusion(s) or other benefit limitations applicable to this service or supply. If there is a discrepancy between this policy and a Member's plan of benefits, the benefits plan will govern. In addition, coverage may be mandated by applicable legal requirements of a State, the Federal government or CMS for Medicare and Medicaid Members. CMS's Coverage Database can be found on the CMS website. The coverage directive(s) and criteria from an existing National Coverage Determination (NCD) or Local Coverage Determination (LCD) will supersede the contents of this MCP and provide the directive for all Medicare members. References included were accurate at the time of policy approval and publication.

## OVERVIEW

Magnetic Resonance Imaging (MRI) is a non-X-ray (non-ionizing radiation) imaging scan that uses a strong magnetic field and radiofrequency waves to produce detailed cross-sectional views of soft tissues, bones, and vascular structures. These cross-sectional images can be reconstructed, rotated, and displayed in many different planes. Imaging can be performed either without (non-enhanced) or with (contrast enhanced) injection of gadolinium containing contrast material into a vein.

Ultrasound has been shown to have similar diagnostic accuracy when compared to MRI and can be considered in lieu of MRI for evaluation of rotator cuff tears, labral injuries, and bicep tendon tears. It is recommended that the ultrasound be completed at a facility competent in performing and interpreting musculoskeletal ultrasound studies. Ultrasound has the benefit of being portable, does not expose the patient to ionizing radiation, and has dynamic imaging capabilities.

In children and adolescents, joint imaging is not necessarily subject to a failed course of conservative therapy. Early intervention may be appropriate.

## COVERAGE POLICY

Shoulder MRI **may be considered medically necessary** when the following criteria are met:

**1. Known Tumor or Mass**

- a. Initial evaluation of a recently diagnosed cancer
- b. Follow up of a known tumor or mass after completion of treatment or with new signs/symptoms
- c. Surveillance of a known tumor or mass according to accepted clinical standards

**OR**

**2. Suspected Tumor or Mass Not Confirmed as Cancer**

- a. Evaluation of an abnormality seen on x-ray or other imaging
- b. Evaluation of an abnormality on physical examination and initial evaluation with x-ray or ultrasound has been completed

**OR**

**3. Evaluation of Known or Suspected Infection**

- a. Suspected osteomyelitis and initial x-ray completed

**OR**

**4. Evaluation of Known or Suspected Fractures**

- a. Suspected fracture and x-ray are non-diagnostic
- b. Evaluation of fracture involving the joint space

OR

**5. Shoulder Pain**

- a. Initial x-ray has been performed and there has been at least 4 weeks of conservative therapy\*

\* Conservative therapy consists of a combination of passive modalities such as rest, ice, activity modification, splinting or use of sling, and active modalities such as physical therapy, a supervised home exercise program, and/or failed injections.

OR

**6. Wrist Pain**

- a. Initial x-ray has been performed and there has been at least 4 weeks of conservative therapy\*
- b. Hemarthrosis – blood in the joint.
- c. Suspected ligament tear with instability on examination or with joint space widening on stress view x-rays
- d. Locked wrist.
- e. For suspected TFCC (triangular fibrocartilage complex) tear
- f. MRI arthrogram

\* Conservative therapy consists of a combination of passive modalities such as rest, ice, activity modification, splinting or bracing, and active modalities such as physical therapy, a supervised home exercise program, and/or failed injections.

OR

**7. Other**

- a. Evaluation of suspected avascular necrosis (AVN) when initial x-ray is non-diagnostic
- b. Evaluation of known or suspected autoimmune disease and x-rays are non-diagnostic and there is consideration to change the treatment regimen. Imaging should be limited to the most symptomatic joint
- c. Evaluation of osteochondral defects or osteochondritis dissecans
- d. Evaluation of an abnormality seen on other imaging and the diagnosis remains uncertain
- e. For evaluation of the brachial plexus

**Pre / Post-Procedural**

- Pre-operative evaluation.
- Post-operative for routine recommended follow up or for potential post-operative complications.
- A repeat study may be needed to help evaluate a patient's progress after treatment procedure intervention or surgery. The reason for the repeat study and that it will affect care must be clear.

**Additional Critical Information**

The above medical necessity recommendations are used to determine the best diagnostic study based on a Member's specific clinical circumstances. The recommendations were developed using evidence-based studies and current accepted clinical practices. Medical necessity will be determined using a combination of these recommendations as well as the Member's individual clinical or social circumstances.

- Tests that will not change treatment plans should not be recommended.
- Same or similar tests recently completed need a specific reason for repeat imaging.

**DOCUMENTATION REQUIREMENTS.** Molina Healthcare reserves the right to require that additional documentation be made available as part of its coverage determination; quality improvement; and fraud; waste and abuse prevention processes. Documentation required may include, but is not limited to, patient records, test results and credentials of the provider ordering or performing a drug or service. Molina Healthcare may deny reimbursement or take additional appropriate action if the documentation provided does not support the initial determination that the drugs or services were medically necessary, not investigational, or experimental, and otherwise within the scope of benefits afforded to the member, and/or the documentation demonstrates a pattern of billing or other practice that is inappropriate or excessive.

## CODING & BILLING INFORMATION

### CPT Codes

CPT	Description
73221	Magnetic resonance (e.g., proton) imaging, any joint of upper extremity; without contrast material(s)
73222	Magnetic resonance (e.g., proton) imaging, any joint of upper extremity; with contrast material(s)
73223	Magnetic resonance (e.g., proton) imaging, any joint of upper extremity; without contrast material(s), followed by contrast material(s) and further sequences
73218	Magnetic resonance (e.g., proton) imaging, upper extremity, other than joint; without contrast material(s)
73219	Magnetic resonance (e.g., proton) imaging, upper extremity, other than joint; with contrast material(s)
73220	Magnetic resonance (e.g., proton) imaging, upper extremity, other than joint; without contrast material(s), followed by contrast material(s) and further sequences

**CODING DISCLAIMER.** Codes listed in this policy are for reference purposes only and may not be all-inclusive. Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement. Listing of a service or device code in this policy does not guarantee coverage. Coverage is determined by the benefit document. Molina adheres to Current Procedural Terminology (CPT®), a registered trademark of the American Medical Association (AMA). All CPT codes and descriptions are copyrighted by the AMA; this information is included for informational purposes only. Providers and facilities are expected to utilize industry standard coding practices for all submissions. When improper billing and coding is not followed, Molina has the right to reject/deny the claim and recover claim payment(s). Due to changing industry practices, Molina reserves the right to revise this policy as needed.

## APPROVAL HISTORY

4/13/2023 Policy reviewed, no changes to criteria.  
 4/13/2022 New policy.

## REFERENCES

### Government Agency

- Centers for Medicare and Medicaid Services (CMS). Medicare coverage database. Accessed February 20, 2023. <http://www.cms.gov/medicare-coverage-database/>.

### Peer Reviewed Publications

- Anderson MW, Greenspan A. Stress fractures. Radiology. 1996 Apr;199(1):1-12. doi: 10.1148/radiology.199.1.8633129. PMID: 8633129.
- Boutry N, Morel M, Flipo RM, et al. Early rheumatoid arthritis: A review of MRI and sonographic findings. AJR Am J Roentgenol. 2007; 189:1502–1509. doi: 10.2214/AJR.07.2548.
- De Coninck T, Ngai SS, Tafur M, et al. Imaging the glenoid labrum and labral tears. Radiographics. 2016 Oct;36(6):1628-1647. doi: 10.1148/rg.2016160020. PMID: 27726737.
- George E, Tsipias S, Wozniak G, et al. MRI of the knee and shoulder performed before radiography. J Am Coll Radiol. 2014 Nov;11(11):1053-8. doi: 10.1016/j.jacr.2014.05.014. PMID: 25086957.
- Haims AH, Moore AE, Schweitzer ME, et al. MRI in the diagnosis of cartilage injury in the wrist. AJR Am J Roentgenol. 2004 May;182(5):1267-70. doi: 10.2214/ajr.182.5.1821267. PMID: 15100130.
- Hayter CL, Gold SL, Potter HG. Magnetic resonance imaging of the wrist: Bone and cartilage injury. J Magn Reson Imaging. 2013 May;37(5):1005-19. doi: 10.1002/jmri.23845. PMID: 23606139.
- Itoi E. Rotator cuff tear: Physical examination and conservative treatment. J Orthop Sci. 2013 Mar;18(2):197-204. doi: 10.1007/s00776-012-0345-2. PMID: 23306597. PMCID: PMC3607722.
- Johnson GW, Cadwallader K, Scheffel SB, et al. Treatment of lateral epicondylitis. Am Fam Physician. 2007; 76(6):843-848. PMID: 17910298.
- Karchevsky M, Schweitzer ME, Morrison WB, et al. MRI findings of septic arthritis and associated osteomyelitis in adults. AJR Am J Roentgenol. 2004 Jan;182(1):119-22. doi: 10.2214/ajr.182.1.1820119. PMID: 14684523.
- Magee T. Usefulness of unenhanced MRI and MR arthrography of shoulder in detection of unstable labral tears. AJR Am J Roentgenol. 2015 Nov;205(5):1056-60. doi: 10.2214/AJR.14.14262. PMID: 26496553.
- Ng AWH, Chu CM, Lo WN, et al. Assessment of capsular laxity in patients with recurrent anterior shoulder dislocation using MRI. AJR Am J Roentgenol. 2009 Jun;192(6):1690-5. doi: 10.2214/AJR.08.1544. PMID: 19457836.
- Pierce JL, Nacey NC, Jones S, et al. Postoperative shoulder imaging: Rotator cuff, labrum, and biceps tendon. Radiographics. 2016 Oct;36(6):1648-1671. doi: 10.1148/rg.2016160023. PMID: 27726742.
- Roy EA, Cheyne I, Andrews G, et al. Beyond the cuff: MR imaging of labroligamentous injuries in the athletic shoulder. Radiology. 2016 Feb;278(2):316-32. doi: 10.1148/radiol.2015150364. PMID: 26789600.

# Molina Clinical Policy

## Shoulder MRI: Policy No. 663

Last Approval: 4/13/2023

Next Review Due By: April 2024



14. Taneja AK, Kattapuram SV, Chang CY, et al. MRI findings of rotator cuff musculotendinous junction injury. *AJR Am J Roentgenol.* 2014 Aug;203(2):406-11. doi: 10.2214/AJR.13.11474. PMID: 25055277.
15. Termaat MF, Raijmakers PGHM, Scholten HJ, et al. The accuracy of diagnostic imaging for the assessment of chronic osteomyelitis: A systematic review and meta-analysis. *J Bone Joint Surg Am.* 2005 Nov;87(11):2464-71. doi: 10.2106/JBJS.D.02691. PMID: 16264122.
16. Tuite M, Small KM. Imaging evaluation of nonacute shoulder pain. *AJR Am J Roentgenol.* 2017 Sep;209(3):525-533. doi: 10.2214/AJR.17.18085. PMID: 28537759.

### National and Specialty Organizations

1. Expert Panel on Musculoskeletal Imaging, Torabi M, Lenchik L, et al. ACR appropriateness criteria ® acute hand and wrist trauma. *J Am Coll Radiol.* 2019 May;16(5S):S7-S17. doi: 10.1016/j.jacr.2019.02.029. Accessed February 20, 2023Expert Panel on Musculoskeletal Imaging, Kransdorf MJ, Murphey MD, et al. ACR appropriateness criteria ® soft-tissue masses. *J Am Coll Radiol.* 2018 May;15(5S):S189-S197. doi: 10.1016/j.jacr.2018.03.012. Accessed February 20, 2023.
2. Pedowitz RA, Yamaguchi K, American Academy of Orthopaedic Surgeons, et al. Optimizing the management of rotator cuff problems. *J Am Acad Orthop Surg.* 2011 Jun;19(6):368-79. doi: 10.5435/00124635-201106000-00007. Accessed February 20, 2023.
3. Roberts CC, Kransdorf MJ, Beaman FD, et al. ACR appropriateness criteria follow-up of malignant or aggressive musculoskeletal tumors. *J Am Coll Radiol.* 2016 Apr;13(4):389-400. doi: 10.1016/j.jacr.2015.12.019. Accessed February 20, 2023.
4. Wise JN, Daffner RH, Weissman BN, et al. ACR appropriateness criteria on acute shoulder pain. *J Am Coll Radiol.* 2011 Sep;8(9):602-9. doi: 10.1016/j.jacr.2011.05.008. Accessed February 20, 2023.