

# Molina Clinical Policy

## Gastric Electrical Stimulation: Policy No. 414

Last Approval: 10/09/2024

Next Review Due By: October 2025



### DISCLAIMER

This Molina Clinical Policy (MCP) is intended to facilitate the Utilization Management process. Policies are not a supplementation or recommendation for treatment; Providers are solely responsible for the diagnosis, treatment, and clinical recommendations for the Member. It expresses Molina's determination as to whether certain services or supplies are medically necessary, experimental, investigational, or cosmetic for purposes of determining appropriateness of payment. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that this service or supply is covered (e.g., will be paid for by Molina) for a particular Member. The Member's benefit plan determines coverage – each benefit plan defines which services are covered, which are excluded, and which are subject to dollar caps or other limits. Members and their Providers will need to consult the Member's benefit plan to determine if there are any exclusion(s) or other benefit limitations applicable to this service or supply. If there is a discrepancy between this policy and a Member's plan of benefits, the benefits plan will govern. In addition, coverage may be mandated by applicable legal requirements of a State, the Federal government or CMS for Medicare and Medicaid Members. CMS's Coverage Database can be found on the CMS website. The coverage directive(s) and criteria from an existing National Coverage Determination (NCD) or Local Coverage Determination (LCD) will supersede the contents of this MCP and provide the directive for all Medicare members. References included were accurate at the time of policy approval and publication.

### OVERVIEW

**Gastroparesis**, or delayed gastric emptying, is a condition in which the stomach fails to empty its contents at a normal rate, which may lead to symptoms including bloating, discomfort, nausea, vomiting, and early satiety. In severe cases, these symptoms can lead to weight loss, dehydration, electrolyte disturbances, and/or malnutrition. Conservative treatments for gastroparesis include medication, glycemic control in the case of diabetes, and dietary modification. Supportive measures such as intravenous hydration or nutritive support may be needed. When conservative management fails to relieve symptoms, the condition may be referred to as refractory (Hasler 2023).

**Gastric electrical stimulation (GES)** is a treatment for refractory gastroparesis which involves implantation of electrodes into the gastric muscles at the greater curvature of the stomach. The electrodes are attached to a small, battery-powered pulse generator placed within a subcutaneous pocket in the abdominal area and programmed remotely to optimize therapy for the individual patient. The procedure is usually performed laparoscopically, although an open approach is available when laparoscopic placement is not an option. Once placed, the generator is programmed to send electric pulses to the stomach muscle wall (Shanker et al. 2021; Hasler 2023).

The mechanism of action of GES in the treatment of gastroparesis is not fully understood. Initially, it was thought that the electrical impulses may improve the rate of gastric emptying, however studies have not shown this to be consistently true. Reduction of symptoms in patients with refractory gastroparesis may be due to neurostimulation travelling to the brain from the stomach via the vagal afferents and affecting the nausea and vomiting centers of the brain. It has also been noted that patients treated with GES appear to have an increased gastric accommodation, which could theoretically lead to symptom relief (Shanker et al. 2021; Hasler 2023).

#### **Regulatory Status**

The GES system, now the Enterra™ Therapy System (Medtronic, Inc.), received approval from the United States Food and Drug Administration (FDA) under the Humanitarian Device Exemption in 2000 as indicated for treatment of chronic, intractable nausea and vomiting secondary to gastroparesis of diabetic or idiopathic etiology (FDA 2000). Temporary GES has been investigated as a less invasive trial to determine the potential efficacy of pacing prior to more invasive long-term placement as described above. In temporary GES trials, cardiac pacers placed endoscopically are connected to an external GES device. This is an off-label use, as the only GES configuration currently approved by the FDA uses pacing wires implanted intramuscularly in the antrum of the stomach.

On October 20, 2023, the Enterra II® System received FDA supplemental approval for Magnetic Resonance conditional use. The MR Conditional label means that patients who receive or have received the Enterra II System in the past can safely undergo Magnetic Resonance Imaging (MRI) examinations of the head or upper/lower extremity under certain conditions.

Gastric Electrical Stimulation (GES) therapy has been proposed as a treatment for patients with obesity. The goal of GES in treating obesity is to cause early satiety and reduce appetite causing subsequent weight loss. The exact mechanisms that result in changes in eating and behavior are uncertain. There are no GES devices approved by the FDA for the treatment of obesity.

# Molina Clinical Policy

## Gastric Electrical Stimulation: Policy No. 414

Last Approval: 10/09/2024

Next Review Due By: October 2025



### COVERAGE POLICY

Gastric electrical stimulation for the treatment of refractory gastroparesis **may be considered medically necessary** when **ALL** the following criteria are met:

1. Diagnosis of gastroparesis with diabetic or idiopathic etiology
2. Delayed gastric emptying, defined by > 60% retention at two hours or > 10% retention at four hours, as measured by standardized gastric emptying study (e.g., scintigraphy)
3. Age 18 or greater
4. Symptoms persist for  $\geq 1$  year despite medical management including:
  - a. Dietary modification
  - b. Refractory to, intolerant to, or has contraindications to the use of at least one drug from **both** of the following classes:
    - i. Antiemetic medication(s) (e.g., antihistamines, serotonin receptor antagonists, and dopamine receptor antagonists) such as granisetron or ondansetron
    - ii. Prokinetic medication(s) (e.g., cholinergic agonists, motilin receptor agonists, and dopamine receptor antagonists) such as metoclopramide

### Limitations and Exclusions

Gastric electrical stimulation is **considered not medically necessary** and may not be authorized for any of the following:

1. Treatment of conditions other than chronic refractory gastroparesis including obesity
2. Use in pregnant individuals
3. Use in individuals under 18
4. Temporary GES in which leads are placed endoscopically

**DOCUMENTATION REQUIREMENTS.** Molina Healthcare reserves the right to require that additional documentation be made available as part of its coverage determination; quality improvement; and fraud; waste and abuse prevention processes. Documentation required may include, but is not limited to, patient records, test results and credentials of the provider ordering or performing a drug or service. Molina Healthcare may deny reimbursement or take additional appropriate action if the documentation provided does not support the initial determination that the drugs or services were medically necessary, not investigational, or experimental, and otherwise within the scope of benefits afforded to the member, and/or the documentation demonstrates a pattern of billing or other practice that is inappropriate or excessive.

### SUMMARY OF MEDICAL EVIDENCE

#### Gastroparesis

Evidence in published peer-reviewed medical literature examining the safety and efficacy of permanent GES for the treatment of refractory gastroparesis consists of observational studies, case series, retrospective studies, and a small number of randomized controlled trials. A summary of the most notable studies is below.

#### **Randomized Controlled Trials**

Ducrotte et al. (2020) conducted a large, multicenter, randomized, double-blind, crossover trial (NCT00903799) that studied the efficacy of GES with the Enterra device in patients with chronic (>12 months) refractory vomiting, with or without gastroparesis. Patients (n=172) were then assigned to double-blind groups that received 4 months of stimulation or no stimulation, and then evaluated on the primary outcomes of vomiting score and quality of life, and secondary outcomes of changes in other digestive symptoms, nutritional status, gastric emptying, and glycemic control. During both phases of the crossover, vomiting was reduced in the group with the device on versus with the device off.

## Molina Clinical Policy

### Gastric Electrical Stimulation: Policy No. 414

Last Approval: 10/09/2024

Next Review Due By: October 2025



The reduction in vomiting was statistically significant ( $P < 0.01$ ) when the device was on in patients with delayed gastric emptying, whereas the reduction in vomiting was just above the threshold of statistical significance ( $P = 0.05$ ) for patients with normal gastric emptying. Gastric emptying was not accelerated when the device was on, nor was the device being on associated with increased quality of life.

McCallum et al. (2013) conducted a prospective, multicenter, double-blind, randomized, crossover study evaluating the efficacy and safety of GES in the treatment of chronic vomiting in gastroparesis of idiopathic etiology. Thirty-two subjects were enrolled and had an Enterra GES system implanted and programmed. Stimulation was turned ON for 1.5 months, followed by double-blind randomization to 3-month crossover periods with the device either ON or OFF, followed by unblinded ON stimulation for 4.5 months. The primary outcome evaluated was weekly vomiting frequency (WVF). There was a significant reduction in WVF from baseline in the initial unblinded ON period (61.2%,  $P < 0.001$ ) and at the 1-year follow up (87%,  $P < 0.001$ ). However, the double-blind 3-month periods showed a non-significant reduction in WVF in the ON versus OFF period, raising the question of placebo effect or whether the lack of a washout after the ON period may have allowed for the continued effects of GES to bleed into the OFF period.

McCallum et al. (201) conducted a prospective, multicenter, randomized, controlled, crossover study (NCT00157755) evaluating the effects of GES with the Enterra therapy system on symptoms from diabetic gastroparesis. Inclusion criteria required more than 7 episodes of vomiting per week, gastric emptying confirmed by scintigraphy ( $> 60\%$  retention at 2 hours and  $> 10\%$  at 4 hours), presence of symptoms for 12 months or greater, and refractoriness or intolerance to prokinetic and antiemetic drug classes trialed for at least 1 month unless contra-indicated. Fifty-five patients with refractory diabetic gastroparesis were enrolled and had the Enterra gastric stimulation system implanted. After placement, the stimulator was turned ON for 6 weeks in all patients, followed by randomization to 3-month crossover periods with the device ON or OFF, followed by unblinded ON setting for 4.5 months. There was a significant reduction in weekly vomiting frequency (WVF) of 57% compared to baseline ( $P < 0.001$ ), and at one year remained significant at 67.8% ( $P < 0.001$ ). There was no difference, however, in WVF between patients who had the device ON or OFF during the blinded 3-month crossover period.

Abell et al. (2003) conducted an early randomized, double-blind crossover trial that enrolled 33 patients ages 19 to 65 years with chronic gastroparesis (17 diabetic, 16 idiopathic) to evaluate the effectiveness of GES on medically refractory gastroparesis. Inclusion criteria required more than 7 episodes of vomiting per week, gastric emptying confirmed by scintigraphy ( $>60\%$  retention at 2 hours and  $>10\%$  at 4 hours), presence of symptoms for 12 months or greater, and refractoriness or intolerance to 2 of 3 classes of prokinetic drugs and 2 of 3 classes of antiemetics. After implantation, patients were randomized to stimulation ON or OFF for 1-month periods. The blind was then broken, and all participants were treated with GES and evaluated again at 6 and 12 months. The primary outcomes evaluated were vomiting frequency, preference for ON or OFF, upper gastrointestinal tract symptoms, quality of life, gastric emptying, and adverse events. During the initial double-blind period, subjects reported significantly reduced vomiting in the ON period as opposed to the OFF ( $P < 0.05$ ). Patient preference for the ON period mirrored the vomiting outcome. Unblinded participants reported significantly decreased frequency of vomiting ( $P < 0.05$ ) at 6- and 12-month follow up. Infection or other complication resulted in 5 individuals having their GES system explanted or revised.

#### **Systematic Reviews and Meta-Analyses**

Rajamanuri et al. (2021) performed a systemic review on literature published in the last 10 years. A total of 1924 articles were identified in databases (PubMed, PubMed Central, Medline, Science Direct, and Google Scholar). The articles focused on gastroparesis and its symptoms or GES therapy. After screening, 124 articles met initial inclusion and exclusion criteria – 41 articles were found most relevant. For the review, 12 studies were included (review articles, observational studies, and clinical trials). Overall, the studies demonstrated efficacy of the pacemaker on symptoms of gastroparesis in adults (pediatric population excluded). Additional RCTs (Randomized controlled trials) are needed that provide an analysis of the efficacy of gastric pacemakers in improving symptoms of gastroparesis. Future research on the use of gastric pacemakers in idiopathic and post-surgical gastroparesis. The authors concluded that gastric pacemakers have shown differing effects on symptoms (e.g., nausea, vomiting, abdominal bloating, weight loss, and overall quality of life). Significant weight gain from this therapy was also noted in the studies reviewed. Many studies indicated a significant improvement in QOL and Gastroparesis Cardinal Symptom Index scores however, a few reported no substantial change in the QOL following GES. Meta-analysis and RCTs vs. open-label trials indicated positive results for QOL because of gastric pacing. The authors note that certain parameters also indicated improvement following GES therapy (e.g., reduction in inflammatory markers, greater insulin levels especially in diabetic patients, and reduced hospital stays).

## Molina Clinical Policy

### Gastric Electrical Stimulation: Policy No. 414

Last Approval: 10/09/2024

Next Review Due By: October 2025



Levinthal and Bielefeldt (2017) conducted a systematic review and meta-analysis of evidence on the use of GES for treatment of gastroparesis and found that total symptom severity scores decreased significantly in open label studies (n = 16), whereas in the studies in which patients were randomized to periods with or without GES there was no significant difference in total symptom severity scores between the two periods. The analysis also showed there were significant differences in baseline total symptoms severity scores among the available studies which significantly impacted the scores during treatment, calling into question whether accurate conclusions can be drawn from the body of evidence.

#### **Non-Randomized Studies, Retrospective Reviews, and Other Evidence**

Gourcerol et al. (2023) completed an observational study comparing gastric electrical stimulation (GES) to gastric peroral endoscopic myotomy (G-POEM) in patients with nausea and vomiting from gastroparesis. Sixty-four patients with medically refractory gastroparesis with nausea and vomiting were included in the study. Data was aggregated from the Enterra cohort and the G-POEM cohort. Both independent open-labeled cohorts followed patients for 24 months. Of the 64 patients included in the study, (n=34) were treated with GES and (n=30) were treated with G-POEM. The authors defined the clinical response as a decrease of  $\geq 1$  point in the nausea and vomiting score and the absence of a switch from one to the other technique before the end of the 24 months. Clinical response was noted in 21/34 (61.7%) in patients treated with GES and in 21/30 (70%);  $p=0.60$ ) patients treated with G-POEM. At the 24 month follow up, mean scores of nausea and vomiting subscale decreased in the GES cohort (from 3.0 to 1.6;  $p=0.001$ ) as well at the G-POEM cohort (from 2.6 to 1.2;  $p=0.001$ ). However, when adjusted from baseline (0.28 [-0.77; 0.19];  $p=0.24$ ) there was no difference between the groups. At 24 months the authors did not see a significant difference in the efficacy of GES and G-POEM with medically refractive gastroparesis with predominant nausea and vomiting. Both cohorts had improved symptomatic and quality of life scores at 24 months. The authors noted several limitations to the study. Patients treated with GES had higher nausea and vomiting sub scores and worse quality of life at baseline as compared to the patients treated with G-POEM, however other characteristics were comparable. There was lack of randomization and blinding, a small sample size and different scales were used between the two cohorts. At the conclusion of the study, it remains unknown if GES or G-POEM is more effective for gastroparesis with refractory nausea and vomiting.

Heckert et al. (2016) conducted a single-center prospective study on the effectiveness of GES with the Enterra therapy system for treatment of refractory gastroparesis. A total of 151 patients with refractory gastroparesis (72 diabetic, 73 idiopathic, 6 other) had the Enterra system implanted. The primary outcome was the Clinical Patient Grading Assessment Scale used to assess response based on a patient completed questioner. Of the 138 patients who completed follow up ( $17 \pm 11$  months after implantation), symptoms were improved in 75% of patients with 43% being at least moderately improved.

A prospective study was conducted by Gourcerol et al. (2012) to evaluate the long-term impact of GES on nausea and vomiting in patients with chronic, intractable nausea and vomiting. The study included patients with both delayed and normal gastric emptying. Thirty-one patients were enrolled in implanted with a GES system. Evaluations were conducted at baseline, 6 months, and 5 years after implantation. Four patients were lost to follow-up, 6 had the devices explanted due to lack of improvement, and 1 patient died. Out of the 20 patients evaluated over 5 years, the quality-of-life score showed 27% improvement ( $p < 0.01$ ), including nausea (62%;  $p < 0.01$ ), vomiting (111%;  $p = 0.03$ ), satiety (158%;  $p < 0.01$ ), bloating (67%;  $p < 0.01$ ) and epigastric pain (43%;  $p = 0.03$ ). Patients with delayed gastric emptying and normal gastric emptying before surgery had an improvement rate of 60% and 50%, respectively. Notably, the 6 patients who had the devices explanted due to lack of improvement were not included in the calculations.

#### **Obesity**

There is insufficient published evidence to support the efficacy and safety of GES therapy for promoting weight loss among patients with morbid obesity. Small clinical trials have reported positive outcomes in weight loss and maintenance of weight loss along with minimal complications. The most notable trial, the SHAPE trial, evaluated the difference in the percentage of excess weight loss (EWL) between the control and treatment groups and found EWL was the same for both groups. Shikora et al. (2009) conducted a prospective, randomized, placebo-controlled, double-blind, multicenter study (NCT00200083) called The SHAPE trial compared implantable gastric stimulation therapy with a standard diet and behavioral therapy regimen in 190 participants with obesity by evaluating the difference in the percentage of EWL between the control and treatment groups. All patients underwent implantation with the implantable gastric stimulator and were randomized to 1 of 2 treatment groups: the control group (stimulation off) or treatment group (stimulation on). The patients were evaluated monthly. All individuals who enrolled in this study agreed to consume a diet with a 500-kcal/d deficit and to participate in monthly support group meetings. The procedure resulted



# Molina Clinical Policy

## Gastric Electrical Stimulation: Policy No. 414

Last Approval: 10/09/2024

Next Review Due By: October 2025



in no deaths and a low complication rate. The primary endpoint of a difference in weight loss between the treatment and control groups was not met. The control group lost 11.7% +/- 16.9% of excess weight and the treatment group lost 11.8% +/- 17.6% (P = .717) according to an intent-to-treat analysis. The authors noted that although implantable gastric stimulation as a surgical option for the treatment of morbid obesity is a less complex procedure than current bariatric operations, the results of the present study do not support its application.

### National and Specialty Organizations

In 2022 the **American College of Gastroenterology** (ACG) updated the 2013 ACG guideline on gastroparesis. The ACG presents this guideline as the official practice recommendations of the American College of Gastroenterology. Recommendation 15 states: "Gastric electric stimulation (GES) may be considered for control of GP symptoms as a humanitarian use device (HUD) (conditional recommendation, low quality of evidence)".(Camilleri et al. 2022).

The **National Institute of Diabetes and Digestive and Kidney Diseases** (2018) states that GES may be effective for some patients whose nausea and vomiting do not improve with dietary changes or medications.

The **National Institute for Health and Care Excellence** (2014) published guidance on gastro-electrical stimulation for gastroparesis noting that current evidence on efficacy and safety supports the use of GES as a treatment for chronic, intractable nausea and vomiting due to gastroparesis.

## CODING & BILLING INFORMATION

### CPT (Current Procedural Terminology)

Code	Description
43647	Laparoscopy, surgical; implantation or replacement of gastric neurostimulator electrodes, antrum
43648	Laparoscopy, surgical; revision or removal of gastric neurostimulator electrodes, antrum
43881	Implantation or replacement of gastric neurostimulator electrodes, antrum, open
43882	Revision or removal of gastric neurostimulator electrodes, antrum, open
64590	Insertion or replacement of peripheral, sacral, or gastric neurostimulator pulse generator or receiver, requiring pocket creation and connection between electrode array and pulse generator or receiver
64595	Revision or removal of peripheral, sacral, or gastric neurostimulator pulse generator or receiver, with detachable connection to electrode array
95980	Electronic analysis of implanted neurostimulator pulse generator system (e.g., rate, pulse amplitude and duration, configuration of wave form, battery status, electrode selectability, output modulation, cycling, impedance and patient measurements) gastric neurostimulator pulse generator/transmitter; intraoperative, with programming
95981	Electronic analysis of implanted neurostimulator pulse generator system (e.g., rate, pulse amplitude and duration, configuration of wave form, battery status, electrode selectability, output modulation, cycling, impedance and patient measurements) gastric neurostimulator pulse generator/transmitter; subsequent, without reprogramming
95982	Electronic analysis of implanted neurostimulator pulse generator system (e.g., rate, pulse amplitude and duration, configuration of wave form, battery status, electrode selectability, output modulation, cycling, impedance and patient measurements) gastric neurostimulator pulse generator/transmitter; subsequent, with reprogramming

### HCPCS (Healthcare Common Procedure Coding System)

Code	Description
C1767	Generator, neurostimulator (implantable), non-rechargeable
C1778	Lead, neurostimulator (implantable)
L8679	Implantable neurostimulator, pulse generator, any type
L8680	Implantable neurostimulator electrode, each
L8688	Implantable neurostimulator pulse generator, dual array, non-rechargeable, includes extension

**CODING DISCLAIMER.** Codes listed in this policy are for reference purposes only and may not be all-inclusive. Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement. Listing of a service or device code in this policy does not guarantee coverage. Coverage is determined by the benefit document. Molina adheres to Current Procedural Terminology (CPT®), a registered trademark of the American Medical Association (AMA). All CPT codes and descriptions are copyrighted by the AMA; this information is included for

# Molina Clinical Policy

## Gastric Electrical Stimulation: Policy No. 414

Last Approval: 10/09/2024

Next Review Due By: October 2025



informational purposes only. Providers and facilities are expected to utilize industry standard coding practices for all submissions. When improper billing and coding is not followed, Molina has the right to reject/deny the claim and recover claim payment(s). Due to changing industry practices, Molina reserves the right to revise this policy as needed.

### APPROVAL HISTORY

<b>10/09/2024</b>	Policy reviewed, no changes to criteria. Updated Overview, Summary of Evidence, and References.
<b>10/12/2023</b>	Policy reviewed, no changes to criteria. Updated Overview and Summary of Evidence due to retirement of policy MCP 243.
<b>06/14/2023</b>	Policy reviewed, no changes to criteria. Updated Summary of Medical Evidence and Reference sections.
<b>06/08/2022</b>	New policy. IRO peer reviewed in May 2022 by a board-certified physician practicing in Gastroenterology.

### REFERENCES

1. Abell T, McCallum R, Hocking M, et al. Gastric electrical stimulation for medically refractory gastroparesis. *Gastroenterology*. 2003 Aug;125(2):421-8. doi: 10.1016/s0016-5085(03)00878-3. PMID: 12891544.
2. Camilleri M, Kuo B, Nguyen L, et al. ACG Clinical Guideline: Gastroparesis. *Am J Gastroenterol*. 2022 Aug 1;117(8):1197-1220. doi: 10.14309/ajg.0000000000001874. Epub 2022 Jun 3. PMID: 35926490; PMCID: PMC9373497.
3. Centers for Medicare and Medicaid Services (CMS). Medicare coverage database. Accessed September 23, 2024. <https://www.cms.gov/medicare-coverage-database/search.aspx>.
4. Ducrotte P, Coffin B, Bonaz B, et al.; ENTERRA Research Group. Gastric Electrical stimulation reduces refractory vomiting in a randomized crossover trial. *Gastroenterology*. 2020 Feb;158(3):506-514.e2. doi: 10.1053/j.gastro.2019.10.018. Epub 2019 Oct 21. PMID: 31647902.
5. Gourcerol G, Gonzalez JM, Bonaz B, et al. Gastric electrical stimulation versus per-oral pyloromyotomy for the treatment of nausea and vomiting associated with gastroparesis: An observational study of two cohorts. *Neurogastroenterol Motil*. 2023 Jul;35(7):e14565. doi: 10.1111/nmo.14565. Epub 2023 Mar 24. PMID: 36961085.
6. Gourcerol G, Huet E, Vandaele N, et al. Long term efficacy of gastric electrical stimulation in intractable nausea and vomiting. *Dig Liver Dis*. 2012 Jul;44(7):563-8. doi: 10.1016/j.dld.2012.01.013. Epub 2012 Mar 3. PMID: 22387288.
7. Hasler WL. Electrical stimulation for gastroparesis. Updated September 12, 2023. Accessed September 20, 2024. <http://www.uptodate.com>.
8. Hayes. Gastric electrical stimulation for gastroparesis Health technology assessment. Published October 26, 2018. Updated December 7, 2022. Accessed September 20, 2024. <https://evidence.hayesinc.com/>.
9. Heckert J, Sankineni A, Hughes WB, et al. Gastric electric stimulation for refractory gastroparesis: A prospective analysis of 151 patients at a single center. *Dig Dis Sci*. 2016 Jan;61(1):168-75. doi: 10.1007/s10620-015-3837-z. Epub 2015 Aug 18. PMID: 26280084.
10. Levinthal DJ, Bielefeldt K. Systematic review and meta-analysis: gastric electrical stimulation for gastroparesis. *Auton Neurosci*. 2017 Jan;202:45-55. doi: 10.1016/j.autneu.2016.03.004. Epub 2016 Apr 11. PMID: 27085627.
11. McCallum RW, Sarosiek I, Parkman HP, et al. Gastric electrical stimulation with Enterra therapy improves symptoms of idiopathic gastroparesis. *Neurogastroenterol Motil*. 2013 Oct;25(10):815-e636. doi: 10.1111/nmo.12185. Epub 2013 Jul 29. PMID: 23895180; PMCID: PMC4274014.
12. McCallum RW, Snape W, Brody F, et al. Gastric electrical stimulation with Enterra therapy improves symptoms from diabetic gastroparesis in a prospective study. *Clin Gastroenterol Hepatol*. 2010 Nov;8(11):947-54; quiz e116. doi: 10.1016/j.cgh.2010.05.020. Epub 2010 Jun 9. PMID: 20538073.
13. National Institute for Health and Care Excellence (NICE). Gastroelectrical stimulation for gastroparesis: [IPG489]. Published May 28, 2014. Accessed September 19, 2024. <https://www.nice.org.uk/guidance/ipg489>.
14. National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK). Gastroparesis. Published January 2018. Accessed September 23, 2024. <https://www.niddk.nih.gov/health-information/digestive-diseases/gastroparesis/treatment>.
15. Rajamanuri M, Mannava SM, Chhabra J, et al. A systematic review of the therapeutic role of gastric pacemakers in adults with gastroparesis. *Cureus [Internet]*. 2021 Sep 21 [cited 2023 May 22];13(9):e18152. doi: 10.7759/cureus.18152. PMID: 34584813; PMCID: PMC8457417.
16. Shanker A, Bashashati M, Rezaie A. Gastric electrical stimulation for treatment of refractory gastroparesis: The current approach to management. *Curr Gastroenterol Rep*. 2021 Jan 22;23(2):2. doi: 10.1007/s11894-020-00803-0. PMID: 33483775; PMCID: PMC7822763.
17. Shikora SA, Bergenstal R, Bessler M, et al. Implantable gastric stimulation for the treatment of clinically severe obesity: Results of the SHAPE trial. *Surg Obes Relat Dis*. 2009 Jan-Feb;5(1):31-7. doi: 10.1016/j.soard.2008.09.012. Epub 2008 Oct 1. PMID: 19071066.
18. United States Food and Drug Administration (FDA). Humanitarian device exemption (HDE) database (product code LNQ, docket number 00M-1451). Gastric electrical stimulation (GES) sys. Issued March 31, 2000. Accessed September 23, 2024. <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfhde/hde.cfm?id=h990014>